

Northern Idaho Fertilizer Guide

Lentils

by Robert L. Mahler

These fertilizer guidelines were developed by the University of Idaho and Washington State University based on relationships between soil tests and crop yield responses. The guidelines are based on research results and are designed to produce above-average yields if other factors are not limiting production. Thus, this fertilizer guide assumes the use of sound management practices.

The suggested fertilizer rates will be accurate for your field provided (1) your soil sample was properly taken and represents the areas to be cropped and (2) the crop and fertilizer history you supply is complete and accurate. For help in obtaining a proper soil sample, refer to University of Idaho Bulletin 704, *Soil Sampling*, or consult the extension educator in your county.

Nitrogen

Lentils are legumes that can obtain or “fix” a portion of the nitrogen (N) they require from the atmosphere. The fixing is done by bacteria (*Rhizobium leguminosarum*) that form nodules on the roots of lentils. These bacteria are present in adequate amounts in most northern Idaho soils.

Seed should be inoculated with this bacterium when (1) lentils have not been grown in the field for 5 or more years before planting or (2) the soil pH is less than 5.7. Lentils are most commonly inoculated with *Rhizobium* in a peat-based carrier using a custom inoculation, seed-applied system.

Follow the inoculum manufacturer’s recommendation for the inoculum rate per bushel of seed. If custom seed inoculation is not available, a peat-based carrier can be used with the planter box, seed-applied system. For additional information on inoculation and methods of

inoculum application, see University of Idaho CIS 838, *Inoculation of Legumes in Idaho*.

In addition to fixed N, the soil often supplies some residual N to lentils as well as N from decomposition of organic matter. Thus, N applications on lentils in most cases have not been profitable. However, lentils need to obtain some N from the soil early in their growth, before effective N-fixing nodules have formed.

Phosphorus

Phosphorus (P) needs can be determined effectively with a soil test (Table 1). Incorporate P into the seedbed by whatever method is most convenient. Acceptable methods include (1) broadcast and plow-down or disk-in, (2) band, and (3) drill with seed. Do not allow direct contact between the seed and any fertilizer containing more than P. Germinating lentils are extremely sensitive to salts contained in fertilizer N, K, and S. If heavy P applications are required to correct nutrient deficiencies, apply fertilizer before or during seedbed preparation.

Table 1. Phosphorus fertilizer rates for lentils based on a soil test.

Soil test P (0 to 12 inches) ¹			Application rate ²	
NaOAc	Bray I	NaHCO ₃	P ₂ O ₅	P
(ppm)	(ppm)	(ppm)	(lb/acre)	(lb/acre)
0 to 2	0 to 20	0 to 8	50	22
2 to 3	20 to 30	8 to 10	30	13
3 to 4	30 to 40	10 to 12	10 ³	5
over 4	over 40	over 12	0	0

¹Soil test P can be determined by three different procedures: sodium acetate (NaOAc), Bray I method, or by sodium bicarbonate (NaHCO₃). Sodium bicarbonate should not be used on soils with pH values less than 6.2. Use the column indicated by your soil test report.

²P₂O₅ × 0.44 = P, or P × 2.29 = P₂O₅.

³Under reduced tillage apply up to 20 lb P₂O₅ per acre on soils testing in excess of 4 ppm P (NaOAc soil test).

Potassium

Potassium (K) needs can be determined with a soil test (Table 2). Incorporate K into the seedbed by whatever method is convenient. Acceptable methods include (1) broadcast and plow-down or disk-in, (2) band, and (3) drill with seed. Do not allow direct contact between the seed and the fertilizer because peas and lentils are sensitive to salts during germination. If heavy K applications are required to correct nutrient deficiencies, apply fertilizer before or during seedbed preparation.

Table 2. Potassium fertilizer rates for lentils based on a soil test.

Soil test K (0 to 12 inches) ¹ (ppm)	Application rate ²	
	K ₂ O (lb/acre)	K (lb/acre)
0 to 50	70	60
50 to 75	40	34
more than 75	0	0

¹ Sodium acetate-extractable K in the 0- to 12-inch depth.

² K₂O x 0.83 = K, or K x 1.20 = K₂O.

Sulfur

Without adequate sulfur (S), lentil plants are unable to fix enough atmospheric N to meet their needs. Consequently, soils testing at less than 10 ppm (SO₄-S) should receive 15 pounds of S per acre. Avoid using granular elemental S on lentils because this form of S becomes available only slowly. Elemental S also greatly reduces soil pH. Sulfur needs of lentils based on a soil test are shown in Table 3.

Table 3. Sulfur fertilizer needs of lentils based on a soil test.

Soil test S (0 to 12 inches)		S application rate
(ppm SO ₄ -S)	(ppm S)	(lb/acre)
0 to 10	0 to 4	15
over 10	over 4	0

Micronutrients

Boron—Lentils grown in northern Idaho respond to boron (B) applications. Boron need can be determined by a soil test. Soils testing at less than 0.5 ppm B should receive 1 pound of B per acre. Boron can be toxic at excessive rates or when concentrated near seedlings. Boron fertilizer should always be broadcast, never banded. For more information on B and specific fertilizer materials, refer to University of Idaho CIS 1085, *Essential Plant Micronutrients: Boron in Idaho*.

Molybdenum—Lentils grown in northern Idaho also respond to molybdenum (Mo). Because Mo is present in soil in only small amounts, soil Mo analysis is not commercially available. Consequently, base Mo fertilizer applications on cropping history and soil pH. Apply Mo as a seed treatment on lentils at the rate of 1/8 to 1/2

ounce Mo per acre when (1) the soil pH is less than 5.7 or (2) every third time lentils are grown in a field. Do not exceed 1/2 ounce Mo per acre because at higher rates the N-fixing bacteria may die. For more information on Mo, refer to University of Idaho CIS 1087, *Essential Plant and Animal Micronutrients: Molybdenum in Idaho*.

Zinc—Response of lentils to zinc (Zn) applications is extremely rare. Zinc applications of 5 pounds per acre should be considered only where Zn soil test levels are less than 0.6 ppm. For more information on Zinc, refer to University of Idaho CIS 1088, *Essential Plant Micronutrients: Zinc in Idaho*.

Other micronutrients—Lentils have reportedly not responded to applications of chlorine (Cl), copper (Cu), iron (Fe), or manganese (Mn). Therefore, applications of these materials in northern Idaho are unnecessary.

Lime—Consider lime applications of 1 ton per acre on fields with a pH of 5.3 and below. Reduced lentil yields may occur at soil pH of 5.4 or lower. However, the yield response from liming at pH values near 5.4 may not always be economical. For additional information on the impact of soil pH on lentils, refer to University of Idaho CIS 811, *The Relationship of Soil pH and Crop Yields in Northern Idaho*.

Agronomy/Water quality considerations

- Weeds, insects, diseases, and environmental stress can influence the effectiveness of a fertilizer program and reduce yields.
- Lentils take up residual soil nitrates (nitrates not used by the preceding cereal crops) and therefore reduce the potential for N loss by leaching. Thus, lentils can have a positive impact on groundwater quality.
- Lentils are capable of fixing most of the nitrogen they need from the atmosphere. Soils in northern Idaho generally contain adequate amounts of the soil bacteria (rhizobia) that are responsible for this nitrogen-fixation process. Consequently, inoculation of lentils with rhizobia is not necessary in fields that have a history of pea or lentil production.
- Early planting of lentil varieties is critical for maximum economic yields.
- Using lentils in a crop rotation can reduce disease and weed problems in grain crops.
- Spring-planted lentils have generally been planted in seedbeds having a minimum of straw residue on the soil surface. However, lentils grown under conservation tillage systems with moderate levels of surface residue typically produce similar or higher yields than lentils grown under low-residue, intensive tillage systems. The greatest yield benefits are in relatively dry years.

- To prevent soil compaction, avoid tillage at high soil moisture levels. Also avoid overworking the soil and creating a finely pulverized surface that is vulnerable to erosion and prone to sealing and crusting.
- Avoid planting in poorly drained areas.
- If you need further information on cultural practices contact the extension educator in your county.
- Starter, or pop-up, fertilizers have limited success on lentils. Starter fertilizers have been most effective when soils were cold and root growth could be stimulated by a readily available supply of P.
- Banding fertilizer improves P use efficiency. Consequently, if applying P in a band, cut the recommended fertilizer application rate by 10 to 15 percent.

Further reading

- BUL 704, *Soil Sampling*, \$2.00
 CIS 811, *The Relationship of Soil pH and Crop Yields in Northern Idaho*, 35 cents
 CIS 838, *Inoculation of Legumes in Idaho*, 35 cents
 CIS 1085, *Essential Plant Micronutrients: Boron in Idaho*, \$3.00
 CIS 1087, *Essential Plant and Animal Micronutrients: Molybdenum in Idaho*, \$1.00
 CIS 1088, *Essential Plant Micronutrients: Zinc in Idaho*, \$3.00

To order copies of these or other University of Idaho Extension publications, contact the University of Idaho Extension office in your county or write to Publications, University of Idaho, P.O. Box 442240, Moscow, ID 83844-2240, call (208) 885-7982, email calspubs@uidaho.edu, or go online at <http://info.ag.uidaho.edu>

Northern Idaho fertilizer guides are available online and may be downloaded from <http://info.ag.uidaho.edu/catalog/catalog.html>. Look under Fertilizers and Soils:

- CIS 447, *Alfalfa*
 CIS 453, *Winter Wheat*
 CIS 785, *Winter Rapeseed*
 CIS 788, *Bluegrass Seed*
 CIS 815, *Blueberries, Raspberries, and Strawberries*
 CIS 820, *Grass Seedings for Conservation Programs*
 CIS 826, *Chickpeas*
 CIS 851, *Legume and Legume-Grass Pastures*
 CIS 853, *Grass Pastures*
 CIS 911, *Northern Idaho Lawns*, also available in print for \$1.00
 CIS 920, *Spring Barley*
 CIS 954, *Winter Barley*
 CIS 1012, *Spring Canola*
 CIS 1083, *Lentils*
 CIS 1084, *Spring Peas*
 CIS 1101, *Soft White Spring Wheat*

The author—Robert L. Mahler is a soil scientist in the University of Idaho Department of Plant, Soil and Entomological Sciences, Moscow.



University of Idaho
Extension

Issued in furtherance of cooperative extension work in agriculture and home economics, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Charlotte V. Eberlein, Director of University of Idaho Extension, University of Idaho, Moscow, Idaho 83844. The University of Idaho provides equal opportunity in education and employment on the basis of race, color, religion, national origin, age, gender, disability, or status as a Vietnam-era veteran, as required by state and federal laws.